



Purina®

MEDICATED

MADE TO  
ORDER (MTO)

# Purina® AMP Purinature Turkey Starter Plus - Medicated

Fine Crumble

Purina® Purinature feeds provide complete nutrition to your flock. The Purina Purinature feeds' formulas have an exclusive combination of ingredients, prebiotic vitamins and organic trace minerals to support your flock to maintain good health and produce the well-desired eggs or meat for your family.



## FEATURES & BENEFITS

### Complete nutrition for poults.

Includes Amprolium for the prevention of coccidiosis.

Contains 24% protein to help your 4-week old poult to grow healthy and strong.

Exclusive combination of ingredients, prebiotic and organic trace minerals to support flock performance.

Fortified with vitamins and organic trace minerals to meet the needs of poults in the early stages of life.

Prebiotics and organic trace minerals included to support digestion and nutrient absorption.

Crumble form for easy consumption by the poults.

# Purina® AMP Purinature Turkey Starter Plus - Medicated

FORM	BAG SIZE	LIFE STAGE
Fine Crumble	25kg	0-4 weeks

## GUARANTEED ANALYSIS

CRUDE PROTEIN	MIN.	24.0%	SODIUM	ACTUAL	0.15%
CRUDE FAT	MIN.	3.0%	VITAMIN A	MIN.	10,000 IU/kg
CRUDE FIBRE	MAX.	6.0%	VITAMIN D	MIN.	2,800 IU/kg
CALCIUM	MIN.	1.3%	VITAMIN E	MIN.	30.0 IU/kg
PHOSPHORUS	MIN.	0.9%	SELENIUM		0.30 mg/kg

## FEEDING DIRECTIONS

**Feed this medicated feed as the sole ration from one day of age up to 4 weeks of age.**

- Don't let feeders run empty.
- Clean feeders and waterers regularly.
- Provide fresh, clean water at all times.
- Any ration change should be made gradually.

### IMPORTANT

Follow sound management practices along with disease and sanitation control measures. If there are any questions, contact your veterinarian, Purina dealer or sales representative.

### WARNING

Do not feed to laying birds in production.

### CAUTION

1. Do not use amprolium medicated feeds as treatment for outbreaks of Coccidiosis.
2. If unexpected deaths occur, obtain an accurate diagnosis and follow a veterinarian's or poultry pathologist's recommendations.
3. Directions for use must be carefully followed.
4. Feed is perishable. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, abortion, or death.

### DISCLAIMER OF LIABILITY

Individual results from use of this product may vary based upon differences in customers' programs of management, health, sanitation, breeding, genetics and feeding. Therefore, Cargill Limited and its Dealers do not warrant or guarantee individual results.

